N15 - Equilibrium

Target: I can compare the values of Q and K to determine which direction a reaction will proceed, and can use ICE Tables to perform calculations related to equilibrium.

N15 - Equilibrium

Reaction Quotient – Q and ICE Tables

The Reaction Quotient

If a reaction mixture containing both reactants and products is not at equilibrium, how can we determine in which direction it will proceed?

The answer is to compare the current concentration ratios to the equilibrium constant.

The concentration ratio of the products (raised to the power of their coefficients) to the reactants (raised to the power of their coefficients) is called the **reaction quotient**, **Q**.

Significance of the Reaction Quotient

For the gas phase reaction

$$aA + bB \Leftrightarrow cC + dD$$

the reaction quotient is:

$$Q_{\mathrm{c}} = rac{[\mathrm{C}]^{c}[\mathrm{D}]^{d}}{[\mathrm{A}]^{a}[\mathrm{B}]^{b}} \qquad Q_{\mathrm{p}} = rac{P_{\mathrm{C}}^{c}P_{\mathrm{D}}^{d}}{P_{\mathrm{A}}^{a}P_{\mathrm{B}}^{b}}$$

Set up the same way as K, but the [] values you plug in are not necessarily the same as the ones at equilibrium.

The Reaction Quotient


If K = Q, the system is at equilibrium

If Q > K, there are more products than when at equilibrium, the system shifts to the left, consuming products and forming reactants until equilibrium is achieved

If Q < K, there are more reactants than when at equilibrium, the system shifts to the right, consuming reactants and forming products until equilibrium is achieved


 If Q > K, the reaction will proceed fastest in the reverse direction.

The products will decrease and reactants will increase.




 If Q < K, the reaction will proceed fastest in the forward direction.

The products will increase and reactants will decrease.




 If Q = K, the reaction is already at equilibrium!


The products and reactants will not change.



 If a reaction mixture contains just reactants, then Q = 0, and the reaction will proceed in the forward direction.



 If a reaction mixture contains just products, then $Q = \infty$, and the reaction will proceed in the reverse direction.



Consider this reaction at some temperature:

$$H_2O(g) + CO(g) \leftrightarrows H_2(g) + CO_2(g)$$
 $K = 2.0$

Assume you start with 8 moles of H_2O and 6 moles of CO, in a 1 L container. What is the concentration of H_2O , CO, H_2 , and CO_2 present at equilibrium?

Time for <u>"ICE Tables"</u> – one the most important problem solving technique in the year.

You will use it a lot!

$$H_2O(g) + CO(g) \leftrightarrows H_2(g) + CO_2(g)$$
 $K = 2.0$

Step #1: Write the law of mass action for the rxn

$$2.0 = \frac{[H_2][CO_2]}{[H_2O][CO]}$$

Step #2: We "ICE" the problem, beginning with the Initial concentrations – <u>always check your volume!</u>

$$H_2O(g) + CO(g) \leftrightarrows H_2(g) + CO_2(g)$$

<u>I</u> nitial:	8	6	0	0
Change:	-x	-x	+X	+X
E quilibrium:	8-x	6-x	X	X

Step #3: Plug equilibrium concentrations into the equilibrium expression, and solve for $x H_2O(g) + CO(g) \Rightarrow H_2(g) + CO_2(g)$

Equilibrium: 8	8-x 6-x	X	X
----------------	---------	---	---

$$2.0 = \frac{(x)(x)}{(8-x)(6-x)}$$

$$x=4$$

Step #3: Plug equilibrium concentrations into the equilibrium expression, and solve for $x = H_2O(g) + CO(g) \Rightarrow H_2(g) + CO_2(g)$

Equilibrium:	8-x	6-x	X	X

$$x=4$$

Answer	8-4 = 4M	6-4 = 2 M	4 M	4 M
--------	----------	-----------	-----	-----

Some of These Can Get Really Tricky...

- Quadratic Equations yuuuck
- Perfect squares example: $9 = (5 x)^2$ take square root of both sides... 3 = 5 - x, so x = 2
- Approximations Remember the 5% rule from Honors Chemistry ???? (K<1, x < 5% of initial concentrations)
- Crazy substitutions and rearrangements

But the concept is always the same! You HAVE to practice to see crazy examples!

YouTube Link to Presentation

https://youtu.be/_658doDe_xQ